미래 기후변화 대응 및 불확실성: 앙상블 접근법의 효용

(기후변화 시나리오 전망 및 예측 기후 정보 활용에서)

정유란

응용사업본부, 응용사업팀, 에이펙 기후센터 2017년 8월 31일



Climate Change



- Climate Variation
 - Uncertainty
- Long-term Climate
 - Adaptation
 - Mitigation
- Impact Assessment
 - Sensitivity
 - Vulnerability



Sustainable Development

Climate Change



- Climate Models
 - various climate models (global climate models, regional climate models)

- Ensemble
 - coupled
 - multi-disciplinary

Future Climate Change Scenarios



- Many Scenarios
 - about 27 GCMs

- low resolution
 - over 100~200 km

			Hist	torical			RCP4.5						RCP8.5					
mdlnms	PR	TX	TN	WD	SR	RH	PR	TX	TN	WD	SR	RH	PR	TX	TN	WD	SR	RH
ACCESS1-0	1	1	1				1	1	1				1	1	1			
bcc-csm1-1-m	1	1	1				1	1	1				1	1	1			
bcc-csm1-1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
CanESM2	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
CCSM4	1	1	1				1	1	1				1	1	1			
CESM1-BGC	1	1	1				1	1	1				1	1	1			
CESM1-CAM5	1	1	1				1	1	1				1	1	1			
CMCC-CM	1	1	1				1	1	1				1	1	1			
CMCC-CMS	1	1	1				1	1	1				1	1	1			
CNRM-CM5	1	1	1				1	1	1				1	1	1			
CSIRO-Mk3-6-0	1	1	1				1	1	1				1	1	1			
FGOALS-g2	1	1	1				1	1	1				1	1	1			
FGOALS-s2	1	1	1				1	1	1				1	1	1			
GFDL-CM3	1	1	1				1	1	1				1	1	1			
GFDL-ESM2G	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
GFDL-ESM2M	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
HadGEM2-AO	1	1	1				1	1	1				1	1	1			
HadGEM2-CC	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
HadGEM2-ES	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
inmcm4	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
IPSL-CM5A-LR	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
IPSL-CM5A-MR	1	1	1				1	1	1				1	1	1			
IPSL-CM5B-LR	1	1	1				1	1	1				1	1	1			
MIROC-ESM-CHEM	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
MIROC-ESM	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
MIROC5	1	1	1				1	1	1				1	1	1			
MPI-ESM-LR	1	1	1				1	1	1				1	1	1			
MPI-ESM-MR	1	1	1				1	1	1				1	1	1			
MRI-CGCM3	1	1	1				1	1	1				1	1	1			
NorESM1-M	1	1	1				1	1	1				1	1	1			

Issues / Concerns



- When applied Climate Information of climate modes to Other Models
 - Hydrological Models
 - Agricultural Models ... (crop models or pest/disease models)

Local/Site-Specific Environment




- Site-specific climate: topography, vegetation, land cover
 - distance, elevation,
 - slope effect,
 - heat island effect, cold air drainage effect etc.

Downscaling methods

- temporal


- spatial

대류권 (10-15km)



Uncertainty of Downscaling






Uncertainty of Downscaling



PRCP

TMAX

TMIN




Models vs. Models:

Growth Stage of Soybean



 Growing season: timing of planting, flowering, pod development, and seed filling



Criteria abnormal temperature



Critical growth stages

	Vegetat	ion period	Reproduction period					
Crop	Daga Tamp	Range of	Range of	Egihara Taran				
Crop	Base Temp.	Opt. Temp.	Opt. Temp.	Failure Temp.				
Corboon	10	25-30 (DT),	26-32 (DT),	27				
Soybean	10	19-24 (NT)	22-27 (NT)	37				

(Holmberg, 1973; Gibson and Mullen, 1996)


Crop model simulation:

Crop models



- Crop
 - Cereal: Rice, Maize,
 - Legume: Soybean, Peanut ...
 - Root: Potato, Taro ...

Potential Yield



Climate information:

Observation, ...

Long-term climate change scenarios

Simulation




- Observation climate:
 - Normal year:
 - Normal 8110: no high temperature during growing seasons
 - High temp. : high(extreme) temperature occurred during critical periods


Climate change scenarios:

Results: Observation Climate



Pod number






Days after planting (days)


Jeonju

Results: Climate change scenarios



- Normal vs. High temperature
 - gray & black line: no high temperature
 - red dot line: high temperature occurred





Shift growth stages



- Planting Season
 - mid- May to mid-June (to end-June)



- Changing flowering and grain filling time
 - mid-July and late- August



high temperature frequently occur

Shift growth stages




- Change planting time: easy way to avoid extreme cases, such as high temperature or no rainy days (i.e., conditions in which crops can not grow/cultivate well)
- In the case of Korean soybean, usually plant June 10 in the southern region
 - shift (+/-) 5 days (shorten / delay)
 - for example,
 - shorten May 25, June 5 with based on June 10
 - delay June 15 etc...


Results

Chung et al., 2016: Responses of soybean yield to high temperature stress during growing season: A Case study of the Korean soybean. Korean Journal of Agricultural and Forest Meteorology, 18(4), 188~198 APCC (doi: 10.5532/KJAFM.2016.18.4.188)



Relative change on yield planted at June 10





Sowing date


Sowing date


Results

Chung et al., 2016: Responses of soybean yield to high temperature stress during growing season: A Case study of the Korean soybean. Korean Journal of Agricultural and Forest Meteorology, 18(4), 188~198 APCC (doi: 10.5532/KJAFM.2016.18.4.188)




Relative change on yield planted at June 10





Sowing date

Sowing date



Reproducibility:

Future Climate Change Scenarios



- 9 Climate Change Scenarios (among many scenarios)
 - 8 GCMs
 - 1 RCM
- Simple Quantile Mapping

Model	Origin	Country	Resolution	
KMA-12km	Korea Meteorological Administation	Korea	12.5km x 12.5km	
CanESM2	Canadian Centre for Climate Modeling and Analysis	Canada	2.8° x 2.8°	
GFDL-ESM2G	NOAA/GFDL (Geophysical Fluid Dynamic	USA	2.5° x 2.0°	
GFDL-ESM2M	Laboratory)	USA	2.3 X 2.0	
HadGEM2-CC	Meteorological Office Hadley Center	UK	1.88° x 1.25°	
inmcm4	Institute for Numerical Mathematics	Russia	2° x 1.5°	
IPSL-CM5A-LR	Institute Pierre Simon Laplace	France	3.75° x 1.8°	
MIROC-ESM	Atmosphere and Ocean Research Institute, National			
MIROC-ESM- CHEM	Institute for Environmental Studies, and Japan Agency for Marine-Earth Science and Technology	Japan	2.8° x 2.8°	

Reproducibility of predicted yield

CanESM2

HadGEM2

INMCM4

GFDL-ESM2G

MIROC-ESM-CHEM

KMA125km

IPSL-CM5A-LR

GFDL-ESM2M


OBS-SIM-PYD

MME 9



Reproducibility

- past climate(weather) at the growth stages
- necessary to analyze in more detail whether the downscaled climate data from GCMs during the major crop growth periods reproduced the observed climate accurately enough, and to evaluate the plant responses in the crop model.




Chung U, Kim YU, Seo BS, Seo MC (2017) Evaluation of Variation and Uncertainty in the Potential Yield of Soybeans in South Korea Using Multi-model Ensemble Climate Change Scenarios.

Agrotechnology 6: 158. doi: 10.4172/2168-9881.1000158)

Multi-Model Ensemble (MIME):

Ensemble averaging




Multi Model Ensemble Methods



Ensemble averaging types


Chung U, Kim YU, Seo BS, Seo MC (2017) Evaluation of Variation and Uncertainty in the Potential Yield of Soybeans in South Korea Using Multi-model Ensemble Climate Change Scenarios.



Number of participating



- Number of participating in MME simulation
 - the estimation error decreased as the number of GMCs in ensemble averaging




Forecast information:

Seasonal (1~6 ms) and Sub-Seasonal (7~30 days) forecast information

Additional

Multi-range climate information



Source: http://slideplayer.com/slide/1447561/

Case study: Project Overview

Forecast mixing in daily time-step (S2S and Seasonal) **Hydrology Agriculture** (watershed Risk Assessment & management Framework (Crop modeling) modeling) for Risk Assessment Framework drought forecast information & week~months weights for models Risk components **Drought Drought forecast using** forecast using machine learning and R/S Seasonal forecast(EDI) (SPI, ...) Down-Down-Downscaling scaling scaling S₂S Remote Seasonal **Climate** Sensing **Forecast** Change **Forecast**

Crop Responses (Seasonal)




Crop Model Simulation based on multi-scale climate information

- Assessment change of cropping calendar from Seasonal Forecast information
 - example, change planting(sowing) day, or suggest rotated crop

Crop Responses (Sub-Seasonal: S2S)

 Assessment critical growth stages of crop responses (or yield components) from S2S information

- in cropping season: 1~2 weeks (within 7 ~ 15 days)
- 3 critical periods (e.g., 1~2 weeks after planting, flowering period, or grain filling period during reproductive period, those periods are within 7~15 days)




Year

쌀보리 유전모수 추정 후, 모형 결과 비교

Winter Seasonal Forecast






Jinju (winter barley: 쌀보리)

Winter Seasonal Forecast





Jinju (winter barley: 쌀보리)

Conclusions:


Breeding & Plant Physiology: Genetic information < <u>Big Data</u>
Multi-Disciplinary Researches
< Ensemble


Conclusions - I: Genetic Information



example: TaeGwang (medium and late maturity)

	CSDL	PPSEN	EM-FL	FL-SH	FL-SD	SD-PM	FL-LF	LFMAX	SLAVR	SIZLF	XFRT	WTPSD	SFDUR	SDPDV	PODUR	THRSH	SDPRO	SDLIP
Taegwang	12.70	0.32	18.71	4.92	9.52	35.84	9.56	1.03	350.00	300.00	1.00	0.19	10.10	1.35	29.28	78.00	0.40	0.20






Genetic Information



- Evaluation genetic properties of TaeGwang
 - 16 sites



정유란, 신평, 서명철 (2017): 작물모형의 생물계절 및 잠재수량 예측력 개선 방법 탐색: I. 유전 모수 정보 향상으로 콩의 개화시기 및 잠재수량 예측력 향상이 가능한가? 한국농림기상학회 (in printed).

Conclusions - II: multi-disciplinary study



- multi-sectors combined approach
 - breeding
 - plant physiology (phenology)
 - agro-climatology
 - crop modeling
 - socioeconomics

Bio-economic approach Socio-**Bio-physical** economic Combined under the extreme weather events

Chung et al., 2014: Modeling the effect of a heat wave on maize production in the USA and its implications on food security in the developing world. *Weather and Climate Extreme*, v5-6, 67-77.

(doi: 10.1016/j.wace.2014.07.002)

'4차 산업혁명과 GIS' 온라인 세미나

4차 산업혁명 시대의 성패를 좌우할 '데이터' 특별한 지식 없이도 누구나 데이터 분석 전문가가 될 수 있는 방법을 소개합니다

더 알아보기

'4차 산업혁명' 어떻게 준비하고 계신가요?

모든 것이 데이터화되는 4차 산업혁명 시대의 핵심은 '데이터 활용' 입니다. 막대한 양의 데이터를 어떻게 수집, 분석, 활용하는지가 성패를 좌우하기 때문입니다. 그러므로 경쟁력을 확보하기 위해 서는 데이터 기반 조직으로 변화해야 하며, 데이터 분석가뿐 아니라 조직 내 구성원 모두가 데이터를 이해하고 활용할 수 있어야 합니다.

경청해 주셔서 감사합니다. Q & A

정유란

(uran.chung@gmail.com/uchung@apcc21.org) 응용사업팀,에이펙 기후센터



